



# Profile Of Chest Injury in the Peri-COVID-19 Period: A Single Centre Series

AKASHDEEP SINGH<sup>1</sup>, ASHISH RATHORE<sup>2</sup>, MARIVADA HARI BABU<sup>3</sup>, VIVEK KUMAR SAHU<sup>3</sup>

A  
B  
S  
T  
R  
A  
C  
T

**BACKGROUND:** Chest traumas continue to constitute about 30% of all traumas and contribute to 25-50% of trauma-related deaths. COVID-19 has its primary pathophysiologies in the lung, and can worsen the morbidity and mortality of chest trauma if it occurs concomitantly.

**AIM:** To Examine the profile and outcome of chest trauma across the peri-COVID-19 period.

**MATERIALS AND METHOD:** A retrospective analysis of cases of chest trauma in the peri-COVID-19 period (mid-Nov 2019 to mid-March 2022) at GMC Doda.

**RESULTS:** Eighty-five cases of chest trauma met inclusion criteria within the 28 months period, M:F = 7.5:1, age range 3-80years, mean age (38.60±17.40years) and median age of 37years. There were 54 (63.5%) cases of blunt chest trauma and 31 (36.5%) cases of penetrating chest trauma. Thirty-one (36.5%) patients sustained haemothorax, 9 (10.6%) pneumothorax, 14 (16.5%) haemopneumothorax, 21 (24.7%) rib fractures, and 10 (11.8%) chest wall lacerations. Fifty-one (60%) patients had isolated chest trauma while the remaining 34 (40%) had associated injuries in one or more other organ systems. Closed thoracostomy tube drainage was definitive treatment in 48 (56.5%) patients while emergency thoracotomy was done in 5 (6%) patients. The treatment administered in the remaining 32 (37%) patients included intercostal nerve block for chest pain from rib fractures, wound exploration and wound repair. In the series, 75 (88%) had complete recovery, six (7%) patients left against medical advice, and four (5%) in-hospital mortality was recorded.

**CONCLUSION:** The profile of chest trauma in the peri-COVID-19 period in our centre differed from the pre-COVID-19 years with a higher mortality figure. Management protocol also necessitated certain modifications.

**KEYWORDS:** Chest injury, Peri-COVID-19 era, Haemothorax

## INTRODUCTION

Chest trauma is a significant source of morbidity and mortality<sup>1</sup>, and constitutes about 30% of all traumas and contribute 25-50% of trauma-related deaths. COVID-19 has its primary pathophysiologies in the lung, and can worsen the morbidity and mortality of chest trauma if both occur concomitantly.<sup>2</sup> The invasion of the virus to the lung cells, myocytes and endothelial cells leads to lung injury pathogenesis<sup>3</sup>, hypoxia-related myocyte injury, body immune response, increased damage of myocardial cells, and intestinal and cardiopulmonary changes, accumulation of oxygen free radicals, changes in intracellular pH, accumulation of lactic acid, electrolyte changes and further cellular damage. Symptomatic COVID-19 patients present with symptoms of respiratory infection and manifestation of viral pneumonitis on radiological imaging of the lung. Both symptoms and investigation findings of COVID-19 can mimic chest trauma and make differentiation difficult when both

conditions may coexist in the same patient. However, their differentiation is important because the two conditions have differentiate management protocols.

The diagnosis of COVID-19 is based on the presence of appropriate respiratory features, PCR test results and chest radiological findings. Chest trauma patients also present with the same set of symptoms and signs. Pulmonary contusion is one of the most common lung injuries in chest trauma and usually occurs at the site of the injury, or on the opposite side through countercoup phenomena. The manifestations in the chest CT are patchy airspace opacities and consolidations with non-segmental distribution and subpleural sparing.<sup>4</sup> The radiological manifestation of contusion and lung opacities in chest CT scan of trauma patients with incidental COVID-19 pneumonia is relatively similar. Patchy peripheral consolidation and GGO are common findings among both groups.



© Akashdeep Singh et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY-NC 4.0, which permits unrestricted use, distribution and reproduction in any medium, provided the use is not commercial and the original author(s) and source are cited.  
Submitted on: 12-Oct-2022; Accepted on: 10-Jan-2023

Although subpleural sparing was identified in traumatic patients, which is similar to findings of lung contusion<sup>5</sup>, some studies reported these findings in COVID 19 pneumonia.<sup>6,7</sup> Also, round central opacities are usually not seen in the first images of trauma patients unless complicated by nosocomial infections or fat emboli during admission; this pattern was visualized in our patients with COVID-19. Associated findings of pneumothorax, pneumomediastinum, and rib fractures are seen in chest injury patients. The diagnosis of COVID-19 in chest trauma patient is necessary since the double pathology is associated with significantly higher mortality rate.

This study examines the profile and outcome of chest trauma across the peri-COVID-19 period.

MATERIALS AND METHOD

A retrospective analysis of cases of chest trauma in the peri-COVID-19 period (mid-Nov 2019 to mid-March 2022) at GMC Doda.

Peri COVID-19 period stratification

- Period I: Pre-COVID-19 era (18/11/19 – 18/03/20)
- Period II: COVID-19 era (19/03/20 – 18/03/21)
- Period III: Post COVID-19 era (19/03/21 – 18/03/22)

ASSESSMENT OF PATIENTS<sup>8</sup>

**Clinical assessment:** Administration of checklist for COVID-19 triage (2nd/3rd periods) (figure 1).  
Clinical history and physical examination (figure 2)

Checklist for Covid-19 triage

Name: \_\_\_\_\_ Age: \_\_\_\_\_ Gender [M]/[F] Phone no: \_\_\_\_\_ Address: \_\_\_\_\_

SN	PARAMETER	YES	SCORE
1	Fatigue		1
2	Desaturation (SpO <sub>2</sub> < 94%) on room air		2
3	Cough		1
4	Difficulty in breathing		2
5	Chills with or without rigors		1
6	Muscle pains		1
7	Sore throat		1
8	Diarrhoea		1
9	Vomiting		1
10	Loss of smell		2
11	Loss of taste		2
12	Headache		1
13	Redness and swelling of the eyes		1
<b>Travel and contact</b>			
1	<b>Contact:</b> A contact is a person who experienced any one of the following exposures during the 2 days before and the 14 days after the onset of symptoms of a probable or confirmed case: a. Face-to-face contact with a probable or confirmed case within 1 metre and for more than 15 minutes OR Direct physical contact with a probable or confirmed case. b. Direct care for a patient with probable or confirmed COVID-19 disease without using proper personal protective equipment		5
2	<b>Travel:</b> a. Have you travelled outside the country within the past month b. Have you travelled outside the state within the past month c. Have you visited any of these places within the past month: Lagos, Abuja or the northern states of Nigeria		1
3	<b>Occupations:</b> Do you work in any of these places: Hospital, Bank, Pharmacy shop, Patient medicine shop		1
4	<b>Hospital Admission:</b> Have you been admitted into any hospital in the past 1 month		2
<b>Total score</b>			

If total score >10, invite COVID-19 Case management team to review

Figure 1. Checklist for COVID-19 triage



Figure 2. Clinical photograph of an assault victim who sustained multiple penetrating chest injury

**Radiological assessment:** CXR (figure 3 and 4), e-FAST, Chest USS, Chest CT scan (figure 5)




Figure 3. Chest radiograph of a patient with multiple rib fractures and left tension pneumothorax from blunt chest injury

**Laboratory studies:** PCV, GXM, E/U/Cr, HIV screening, COVID-19 screening, Plasma glucose, Urinalysis



**Figure 4.** Chest radiograph of a patient right haemothorax from penetrating chest injury

**Exclusion and data management:** Patients with incomplete clinical or investigative data were excluded.



**Figure 5.** Chest CT scan of a patient who sustained blunt chest injury

Data analysis was done using SPSS (version 20) and summarized as frequencies and percent.

**RESULTS**

Eighty-five patients met the inclusion criteria for this study amongst those that were admitted for treatment

for chest trauma during the period. They were 75 males and 10 females (M:F = 7.5:1). Of these, young adults aged 20-29years accounted for 28.2%, 30-39years accounted for 18.8%, middle aged adults aged 40-49% accounted for 20% and 50-59years accounted for 8.2%. Other age groups were less frequently affected (table 1). Table 2 shows that there were 54 (63.5%) blunt chest injuries and 31 (36.5%) penetrating chest injuries. Haemothorax occurred in 31 (36.5%) of the cases of chest injury. This was followed by isolated cases of rib fractures in 21 (24.7%) cases, haemopneumothorax in 14 (16.5%) cases, soft tissue injuries in 10 (11.8%) cases and pneumothorax in 9 (10.6%) cases. Table 2 also shows that 31 (40%) patients had associated extra-thoracic injuries.

Variables	Frequency	Percent
<b>Sex</b>		
Male	75	88.2
Female	10	11.8
Total	85	100
<b>Age category</b>		
Less than 10	3	3.6
10-19	4	4.7
20-29	24	28.2
30-39	16	18.8
40-49	17	20.0
50-59	7	8.2
60-69	6	7.1
70-79	6	7.1
80 and above	2	2.4
Total	85	100

**Table 1.** Demographic characteristics of patients with chest injury

Table 3 shows that closed thoracostomy tube drainage was carried in 48 (56.5%) cases of chest injury during the period, while thoracotomy was carried out in only 5 (5.9%) cases. The remaining 37.6% patients were treated with medications, wound care and chest physiotherapy. Seventy-five (88.2%) patients recovered, 7.1% left against medical advice, and mortality rate was 4.7% (table 3).

Table 4 shows the summary of the profile and treatment outcome of patients with injury during the peri COVID-19 period. Fifteen (17.6%), 41 (48.2%) and 29 (34.1%) of the cases occurred in the pre COVID-19, COVID-19 and post COVID-19 periods respectively. Nine (10.6%), 27 (31.8%) and 18 (21.2%) of the blunt chest injury occurred during the three stratified peri COVID-19 periods. And six (7.1%), 14 (16.5%) and 11 (12.9%) of the penetrating chest injuries occurred in the respective peri COVID-19 periods. There was no mortality or discharge against medical advice amongst



Variables	Frequency	Percent
<b>Type of chest injury</b>		
Blunt	54	63.5
Penetrating	31	36.5
Total	85	100
<b>Pathological pattern</b>		
Haemothorax	31	36.5
Pneumothorax	9	10.6
Haemopneumothorax	14	16.5
Rib fracture	21	24.7
Soft tissue injury	10	11.8
Total	85	100
<b>Associated extra-thoracic injury</b>		
Yes	34	40
No	51	60
Total	85	100

**Table 2.** Pathological characteristics of chest injury

patients who sustained chest injury and were treated in our centre with 100% recovery in the pre COVID-19 period. However, during the COVID-19 period, the mortality rate amongst chest injury patient was 4.9% and discharge against medical advice was 7.3% while recovery rate was 87.8%. Also, during the post COVID-19 era, mortality rate amongst our chest injury patients was 7.0%, discharge against medical advice 10.3% and recovery rate was 82.7%. Emergency thoracotomy rate was thrice during COVID-19 period compared to the other periods.

Variables	Frequency	Percent
<b>Treatment</b>		
Closed thoracostomy tube drainage	48	56.5
Thoracotomy	5	5.9
Others	32	37.6
Total	85	100
<b>Outcome</b>		
Recovery	75	88.2
Death	4	4.7
DAMA	6	7.1
Total	85	100

**Table 3.** Treatment and outcome of chest injury

## DISCUSSION

The current study corroborates male preponderance (M:F= 7.5:1) in chest trauma.<sup>2,3,9</sup> This is globally explained by the fact that men engage more in risky activities (legitimate and illegitimate) than women. Majority (67%) of victims were in productive age groups<sup>2,3</sup> (20-29yrs=28%, 30-39yrs=19% and 40-

49yrs=20%) with mean of 38.6±17.4 yrs. The wide age range (youngest 3years old and oldest 80years old) supported non-immunity of any age group to chest trauma.<sup>3</sup>

There were more blunt chest injuries than penetrating chest injuries (63.5%vs36.5%) which is in keeping with global trend and our earlier institutional findings.<sup>10-17</sup> This trend was maintained across the three stratified periods of pre-, peri-, and post-COVID 19 periods. This majorly reflected the aetiopathogenesis and mechanism of chest injury in civilian setting which are dominated by road traffic injuries and falls.

The pathologic entities encountered in our patients' population were same as in earlier studies in the institution.<sup>10,12,13</sup> These included haemothorax (36.5%), pneumothorax (10.6%) and pneumo-haemothorax (16.5%) occurring in association with rib fractures or in isolation, and isolated rib fractures in 24.7%. About 12% of the patients sustained various degrees of chest wall lacerations. Some series may exclude chest wall lacerations from chest injury data. However, the broad definition of chest injury as disruption of tissues of the chest wall and or intrathoracic organs allows inclusion of soft tissue chest wall injuries. Diaphragmatic rupture was not sustained by any patient during this period, as opposed to previously.<sup>13</sup>

This series had associated extra-thoracic injuries in 40% of cases which are known to worsen mortality index in chest trauma.<sup>2,10,13</sup> All recorded mortalities in the series were in patients with associated severe traumatic brain injury and massive haemoperitoneum. Related studies have discovered major associated injuries and worse categories of injury severity scores as some of the determinants of mortality in chest injury.<sup>2,10,13,18</sup>

Management followed the set protocols with additional infection prevention control guidelines re-amplified during the COVID-19 era. Pre-administration of the checklist for COVID-19 triage often resulted in suspiciously high score in many of our patients who presented with low values of peripheral arterial oxygen saturation (SpO<sub>2</sub>). However, no patient tested positive to COVID-19 test. This was explained by the fact that the impact of chest injury manifests pathophysiologically as cardiopulmonary dysfunction same as COVID-19.<sup>1-3,9,19</sup>

About 56% of the patients were successfully treated with closed thoracostomy tube drainage, 5.9%

Variables	Pre-COVID-19(%)	COVID-19 era (%)	Post COVID-19(%)	Total (%)
<b>Types</b>				
Blunt chest injury	9 (10.6)	27 (31.8)	18 (21.2)	54 (63.5)
Penetrating chest injury	6 (7.1)	14 (16.5)	11 (12.9)	31 (36.5)
Total	15 (17.6)	41 (48.2)	29 (34.1)	85 (100)
<b>Treatment</b>				
Tube thoracostomy	6 (7.1%)	23 (27.1)	19 (22.4)	48 (56.5)
Emergency thoracotomy	1 (1.2)	3 (7.3)	1 (1.2)	5 (5.9)
Others	8 (9.4)	15 (17.6)	9 (10.6)	32 (37.6)
Total	15 (17.6)	41 (48.2)	29 (34.1)	85 (100)
<b>Outcome</b>				
Mortality	0 (0.0)	2 (4.9)	2 (7.0)	4 (4.7)
LAMA	0 (0.0)	3 (7.3)	3 (10.3)	6 (7.1)
Recovery rate	15 (100)	36 (87.8)	24 (82.7)	75 (88.2)
Total	15 (100)	41 (100)	29 (100)	85 (100)

**Table 4.** Peri-COVID-19 period distribution of chest injury

underwent thoracotomy as part of their treatment, while the remaining 37.7% of patients were treated non-operatively. These included wound care, chest pain control with intercostal nerve block and/or systemic analgesia and empirical antibiotic and deep vein thrombosis prophylaxis in selected patients. These chest pain control modalities and thoracic epidural analgesia have been in use for chest pain management in the centre.<sup>11,15,16,20</sup> Of the 53 patients that underwent surgical procedures, closed thoracostomy tube drainage was successfully used in 48 (90.6%) of the cases while thoracotomy was needed in only 5 (9.4%). This corroborates global best practice.<sup>12,13</sup> All patients additionally underwent chest physiotherapy.

Although the pre-COVID-19 period reviewed was shorter, across-period analysis seemed to show higher incidences of chest trauma during the COVID-19 and post COVID-19 eras with higher incidence of penetrating chest injuries, higher level of discharge against medical advice and higher mortality rate. Reason attributed to the higher incidence of chest injury in the COVID-19 era include the effects of lockdown which included long periods of social aggregation of persons who could not go to work but also refused to remain at home. Such aggregations, often with substances abuse end in physical fights and assaults. The high mortality rate and discharge against medical advice rate were due to inability to pay for treatment because of the economic hardships occasioned by the lockdown. This is common in Nigeria where health insurance coverage is abysmal.

## CONCLUSION

The profile of chest trauma in the COVID-19 period in our centre differed from the pre-COVID-19 era and was

characterized by higher incidence of discharges against medical advice and a higher mortality figure. Management protocol also necessitated certain modifications including use of checklist for COVID-19 triage aimed at protecting healthcare workers, other patients and the general population against the COVID-19 pandemic.

## REFERENCES

1. Blunt chest trauma: Practice essential, anatomy, pathophysiology, etiology, epidemiology and prognosis
2. Wang Y, Zeng C, Dong L, Liu C, Cai Y, Zhang N, etc. Pulmonary contusion during the COVID 19 pandemic: challenges in diagnosis and treatment. *Surgery Today* 2020;50:1113-6. <https://doi.org/10.1007/s00595-020-02081-9>
3. Sabetian G, Feiz F, Shakibafard A, Fard HA, Sefidbakht S, Jafari SH, etc. Challenges of diagnosis of COVID-19 in trauma patients: A case series. *Trauma* 2021;23(3):218-29
4. Kaewlai R, Avery LL, Asrani AV, et al. Multidetector CT of blunt thoracic trauma. *RadioGraphics* 2008;28:1555-70.
5. Ganie FA, Lone H, Lone GN, et al. Lung contusion: a clinico-pathological entity with unpredictable clinical course. *Bull Emerg Trauma* 2013;1:7-16.
6. Chung M, Bernheim A, Mei X, et al. CT imaging features of 2019 novel coronavirus (2019-nCoV). *Radiology* 2020;295:202-7
7. Yoon SH, Lee KH, Kim JY, et al. Chest radiographic and CT findings of the 2019 novel coronavirus disease (COVID-19): analysis of nine patients treated in Korea. *Korean J Radiol* 2020;21:494-500
8. Sawhney C, Singh Y, Jain K, Sawhney R, Trikha A. Trauma care and COVID-19 pandemic. *J Anaesthesiol Clin Pharmacol* 2020;36:115-20.

9. VanDruff RR, Lee JK. Blunt and penetrating chest trauma with concomitant COVID-19 infections: Two case reports. *Trauma Case Reports* 2021;34:100507
10. Ekpe EE, Eyo C. Determinants of Mortality in Chest Trauma Patients. *Niger J Surg* 2014;20:30-4
11. Ekpe EE, Eyo C. Effect of Analgesia on the Changes in Respiratory Parameters in Blunt Chest Injury with Multiple Rib Fractures. *Ann Afr Med*. 2017;16(3):120-6. [https://doi.org/10.4103/aam.aam\\_73\\_16](https://doi.org/10.4103/aam.aam_73_16).
12. Ekpe EE, Eyo C. Overview of Blunt Chest Injury with Multiple Rib Fractures. *Brit J Med Med Res* 2016;12(8): 1-15.
13. Ekpe EE, Etta O, Akpan AF. Pattern of chest injuries and treatment outcome in a Nigerian Teaching Hospital. *World Journal of Biomedical Research* 2018; 5:32-8
14. Makama JG, Yusuf N, Edaigbini S, Ekpe EE. Thoracostomy tube in trauma surgery: Does position really matter? *Arch Int Surg*. 2018;8:166-70.
15. Ekpe EE, Bassey OO, Etiuma AU, Ikpe MC. Intercostal nerve block proved superior to systemic analgesia in the control of chest pain in patients with traumatic rib fractures. *IOSR Journal of Dental and Medical Science (IOSR-JDMS)* 2018; 17(3):68-74.
16. Ekpe EE, Eyo CS, Bassey OO, Etiuma AU. Analgesia Reverses Abnormal Lung Function Tests in Patients with Blunt Chest Injury. *Journal of Advances in Medicine and Medical Research* 2018;26(11):1-11.
17. Ekpe EE, Uduma F, Umoh V, Ikpe MC, Eyo C, Akpan AF. Effectiveness of small-bore ambulatory pleural drain in the treatment of pneumothorax and haemothorax. *International Journal of Innovative Research in Medical Science* 2019;4(2):163-7.
18. Moon SH, Kim JW, Byun JH, Kim SH, Choi JY, Jang IS, et al. The thorax trauma severity score and the trauma and injury severity score: Do they predict in-hospital mortality in patients with severe thoracic trauma?: A retrospective cohort study. *Medicine (Baltimore)*. 2017;96(42):e8317. <https://doi.org/10.1097/MD.00000000000008317>.
19. Ekpe EE, Umoh IA, Shogade T, Eyo CS, Akpan AF. Review of cardiovascular involvements in COVID-19. *Ibom Medical Journal* 2021;14(4):411-26.
20. Etta. OE, Anyanwu P, Edubio MN, Ekpe EE. Role of thoracic epidural analgesia in the management of blunt chest injuries in a resource-poor setting –a case report. *Ibom Med J*. 2016;9:25-9.

**Cite this article as:**

Singh A, Rathore A, Babu MH, Sahu VK. Profile Of Chest Injury in the Peri-COVID-19 Period: A Single Centre Series. *Int Healthc Res J*. 2023;6(10):OR1-OR9. <https://doi.org/10.26440/IHRJ/0610.01588>

**AUTHOR AFFILIATIONS:** (\*Corresponding Author)

1. Assistant Professor, Department of General Surgery, GMC Doda
2. Senior Resident, Department of General Surgery, GMC Doda
3. DNB Resident, Department of General Surgery, GMC Doda

**Source of support:** Nil, **Conflict of interest:** None declared

**Contact Corresponding Author at:** moishapoo[at]gmail[dot]com